

# PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>PCT 0628</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/NL 97/ 00557</b>	International filing date (day/month/year) <b>07/10/1997</b>	(Earliest) Priority Date (day/month/year) <b>08/10/1996</b>
Applicant <b>UNIVERSITEIT UTRECHT et al.</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. ☐ Certain claims were found unsearchable (see Box I).

2. ☐ Unity of invention is lacking (see Box II).

3. ☐ The international application contains disclosure of a nucleotide and/or amino acid sequence listing and the international search was carried out on the basis of the sequence listing

☐ filed with the international application.

☐ furnished by the applicant separately from the international application,

☐ but not accompanied by a statement to the effect that it did not include matter going beyond the disclosure in the international application as filed.

☐ Transcribed by this Authority

4. With regard to the title, ☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the abstract,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this International Search Report, submit comments to this Authority.


6. The figure of the drawings to be published with the abstract is:

Figure No.            ☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.



# PATENT COOPERATION TREATY

REC'D 20 NOV 1998

WIPO

PCT

## PCT

### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference PCT 0628	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (PCT/IPEA/416)	
International application No. PCT/NL97/00557	International filing date (day/month/year) 07/10/1997	Priority date (day/month/year) 08/10/1996
International Patent Classification (IPC) or national classification and IPC G01N33/554		
Applicant UNIVERSITEIT UTRECHT et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I. ☒ Basis of the report
- II. ☐ Priority
- III. ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV. ☐ Lack of unity of invention
- V. ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI. ☐ Certain documents cited
- VII. ☒ Certain defects in the international application
- VIII. ☐ Certain observations on the international application

Date of submission of the demand 06/05/1998	Date of completion of this report 17. 11. 98
Name and mailing address of the IPEA/  European Patent Office D-80298 Munich Tel. (+49-89) 2399-0. Tx: 523656 epmu d Fax: (+49-89) 2399-4465	Authorized officer Hinchliffe, P Telephone No. (+49-89) 2399-8431 



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/NL97/00557

## I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

### Description, pages:

1-29 as originally filed

### Claims, No.:

1-12 as originally filed

### Drawings, sheets:


1/5-5/5 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/NL97/00557

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes:	Claims
	No:	Claims 1-12
Inventive step (IS)	Yes:	Claims
	No:	Claims 1-12
Industrial applicability (IA)	Yes:	Claims 1-12
	No:	Claims

### 2. Citations and explanations

**see separate sheet**

## VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:

**see separate sheet**






## **SECTION V**

1. The subject matter of the independent claims 1-4 (and dependent claims 5-9) does not fulfill the requirement of Article 33(2) PCT because the said subject matter is not novel.
  - 1.1 The four independent methods all appear to be basically the same method in as much as a peptide expressed on the surface of a replicable display package (such as a phage) is allowed to react with a library of immobilised oligopeptides (i.e. on a solid phase). It appears that the source of the oligopeptides, be they from natural sequence representing antigens or otherwise, can not be truly regarded as being essential to differentiating the independent claims from each other.
  - 1.2 In particular D1 (WO 92 15677) discloses in claim 1 a method for identifying a binding protein expressed on the outer surface of a genetic package (such as a microorganism or phage). The plurality of packages, expressing a variety of peptides on their surfaces, are then brought into contact with the immobilised target (see p.70, section V, "affinity selection of target binding mutants"). The immobilised target is selected from, inter alia, organic macromolecules such as polypeptides (see p.71 I.17-19). Consequently D1 is considered to be novelty destroying for the subject matter of these claims.
  - 1.3 D2 (J.Mol.Biol. (1995), vol.248, p.97-105, De Kruif et al) discloses that a collection of phages, each expressing on its surface a peptide (representing an antibody fragment, such as a single chain Fv) was reacted with an immobilised antigen (see p.99, col.1). Again this method is regarded as being essentially the same as those claimed and consequently the subject matter of claims 1 to 4 is not novel.
2. The subject matter of claims 10,11,12 does not fulfill the requirements of Article 33(2) PCT.
  - 2.1 D2 discloses on p. 99, col.2, that to test for specificity of a phage binding to a particular antigen, the same phage was tested against a panel of irrelevant antigens. This disclosure appears to be prejudicial to the novelty of these claims.

## **SECTION VII**

1. Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the document D1 is not mentioned in the description, nor is this




**WRITTEN OPINION  
SEPARATE SHEET**

---

International application No. PCT/NL97/00557

document identified therein.



## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark  
Office  
(Box PCT)  
Crystal Plaza 2  
Washington, DC 20231  
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

<b>Date of mailing (day/month/year)</b> 29 May 1998 (29.05.98)	
<b>International application No.</b> PCT/NL97/00557	<b>Applicant's or agent's file reference</b> PCT 0628
<b>International filing date (day/month/year)</b> 07 October 1997 (07.10.97)	<b>Priority date (day/month/year)</b> 08 October 1996 (08.10.96)
<b>Applicant</b> LOGTENBERG, Ton et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

06 May 1998 (06.05.98)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO  
34, chemin des Colombettes  
1211 Geneva 20, Switzerland

Authorized officer

L. Panakal




# INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 97/00557

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 15982 A (GENZYME CORP) 15 June 1995 see claims 1-5,65 ---	6
Y	WO 95 31723 A (FRYKBERG LARS ;JACOBSSON KARIN (SE)) 23 November 1995 see abstract ---	7
A	GEYSEN ET AL: "USE OF PEPTIDE SYNTHESIS TO PROBE VIRAL ANTIGENS FOR EPITOPES TO A RESOLUTION OF A SINGLE AMINO ACID" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 81, July 1984, WASHINGTON DC, pages 3998-4002, XP000609021 cited in the application see abstract -----	1,3





## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 97/00557

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9215677 A	17-09-92	US 5223409 A	29-06-93
		JP 7501923 T	02-03-95
		AU 1545692 A	06-10-92
		AU 1578792 A	06-10-92
		AU 1581692 A	06-10-92
		AU 8740491 A	28-04-92
		CA 2105300 A	02-09-92
		CA 2105303 A	02-09-92
		CA 2105304 A	02-09-92
		EP 0575485 A	29-12-93
		EP 0573603 A	15-12-93
		EP 0573611 A	15-12-93
		JP 6510522 T	24-11-94
		JP 7501203 T	09-02-95
		US 5403484 A	04-04-95
		US 5571698 A	05-11-96
		WO 9206191 A	16-04-92
		WO 9215605 A	17-09-92
		WO 9215679 A	17-09-92
		US 5663143 A	02-09-97
WO 9208738 A	29-05-92	AU 662503 B	07-09-95
		AU 9077791 A	11-06-92
		CA 2095735 A	08-05-92
		EP 0556320 A	25-08-93
		JP 6502076 T	10-03-94
WO 9515982 A	15-06-95	AU 1432195 A	27-06-95
		EP 0733070 A	25-09-96
		JP 9506262 T	24-06-97
		CA 2175482 A	15-06-95
WO 9531723 A	23-11-95	AU 2541595 A	05-12-95
		EP 0760100 A	05-03-97



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/NL 97/00557

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 G01N33/554 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 15677 A (PROTEIN ENGINEERING CORP) 17 September 1992 see the whole document	1,3,5-7, 11,12
X	DE KRUIF ET AL.: "Selection and application of human single chain Fv antibody ...." JOURNAL OF MOLECULAR BIOLOGY, vol. 248, no. 1, 21 April 1995, LONDON, pages 97-105, XP000196590 cited in the application see the whole document	1,3,5, 8-12
Y	---	2,4,6,7
Y	WO 92 08738 A (ABBOTT LAB) 29 May 1992 see page 8, line 21 - line 23 --- -/--	2,4

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 December 1997

Date of mailing of the international search report

19.12.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Ceder, 0





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>G01N 33/554, 33/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/15833</b> <b>(43) International Publication Date:</b> 16 April 1998 (16.04.98)
<b>(21) International Application Number:</b> PCT/NL97/00557 <b>(22) International Filing Date:</b> 7 October 1997 (07.10.97) <b>(30) Priority Data:</b> 96202791.8 8 October 1996 (08.10.96) EP <i>(34) Countries for which the regional or international application was filed:</i> NL et al. <b>(71) Applicant (for all designated States except US):</b> UNIVER- SITEIT UTRECHT [NL/NL]; Sorbonnelaan 16, NL-3584 CA Utrecht (NL). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> LOGTENBERG, Ton [NL/NL]; Dorpsstraat 19, NL-3433 CH Vreeswijk (NL). DE KRUIF, Cornelis, Adriaan [NL/NL]; Bemuurde Weerd OZ 17bis, NL-3514 AN Utrecht (NL). <b>(74) Agent:</b> SMULDERS, Th., A., H., J.; Vereenigde Octrooibu- reaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHODS AND MEANS FOR SELECTING PEPTIDES AND PROTEINS HAVING SPECIFIC AFFINITY FOR A TARGET		
<b>(57) Abstract</b>  The invention provides novel methods and means for selecting peptides and for other proteinaceous molecules having specific binding affinity for a target. The binding peptides/proteins are displayed on replicable display packages, preferably phages. The binding peptides are tested for this binding affinity to the target. In one embodiment, the target is provided in the form of peptides displayed on a solid phase, in another the target is presented in a tissue section.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title: Methods and means for selecting peptides and proteins having specific affinity for a target.

This invention relates to the field of proteinaceous biological molecules having specific affinity for a target. More specifically the invention relates to such molecules derived from specific binding peptides such as proteins such  
5 as antibodies, in particular monoclonal antibodies or specific binding derivatives or fragments thereof. In an exemplary embodiment the invention relates to peptides and antibodies which are more or less randomly produced as large collections of different molecules (libraries) expressed on  
10 the surface of a replicable genetic package. From these libraries peptides or antibodies are affinity-selected for binding to the target molecule.

Specific binding affinities between biological molecules or  
15 derivatives thereof have been used throughout applied biotechnology, molecular biology, medical biology, etc. for a very long period of time for many different purposes. Some of the most commonly applied biological molecules having a specific binding affinity are antibodies. Antibodies are  
20 complex proteins usually composed of two light chains and two heavy chains which comprise constant regions and variable regions which variable regions again comprise hypervariable domains, which determine the binding specificity of the antibody (complementarity determining regions, (CDR)).  
25 Antibodies are part of the normal defense system of higher vertebrates. They are produced by cells of the immune system. Until 1975 antibodies were obtained by administering an antigenic substance to an animal (often a rodent) and harvesting the antibodies produced by the animal in its  
30 immune response. This resulted in mixtures of antibodies having many different specificities for different binding sites on the antigenic substance. These mixtures are referred to as polyclonal antisera.

The advent of monoclonal antibodies (Kohler and Milstein (1975) Nature 256:495) represented a significant technical breakthrough with important consequences both scientifically and commercially. Monoclonal antibodies are conventionally made by fusing antibody-secreting plasma cells from an immunized animal with a tumor cell line in order to immortalize the former plasma cells. The immortalized hybridomas are screened for the production of antibodies with the desired specificities. Selected hybridomas are expanded to large numbers and the monoclonal antibodies secreted into the medium are harvested and purified for application. Since the arrival of hybridoma technology, monoclonal antibodies have found broad application in procedures aimed at examining and identifying many antigens in many formats from pregnancy tests to AIDS tests. They are also widely applied in research laboratories for use in studying and isolating a broad variety of molecules and for immunohistochemistry, i.e. identifying certain cells in tissues. For instance, monoclonal antibodies that are reactive with specific antigens expressed on subpopulations of cells can be utilized to detect these cells in tissue sections using immunohistochemical or immunofluorescent approaches. The inherent advantages of monoclonal compared to polyclonal antibodies is that they react only with one specific epitope and that they are standardized from batch to batch and from laboratory to laboratory.

Despite of all the advantages obtainable using monoclonal antibodies, there are a number of limitations associated with hybridoma technology and therefore with the monoclonal antibodies that can be obtained. For instance, the number of hybridomas that can be established and screened for the secretion of a certain antibody specificity is limited. The immunization protocol and the manner in which the animals immune system handles the antigen favors the selection of hybridomas secreting antibodies against immunodominant



epitopes, whereas other specificities will not be obtained because the immunizing antigen or some of its epitopes are not recognized by the animals immune system. In addition, the MoAbs obtained from animals are hardly suitable for  
5 therapeutical use in humans because they evoke an immune respons. Some of these drawbacks may be circumvented by using antibody engineering, aiming at chimeric antibodies of which the variable regions are of animal origin and the constant regions are of human origin (the constant region is  
10 responsible for the bulk of the immune response). However these chimeras still evoke an immune response. Another possibility to avoid the immune response is a technique called CDR-grafting, whereby only the CDR's are of animal origin, which CDR's are inserted in a human antibody  
15 framework. These humanized antibodies are rather difficult to make and often loose some of their binding affinity.

Another way of modifying antibodies is trying to obtain the smallest fragment that will still specifically bind the  
20 antigen (or to be more precise the epitope) with a significant affinity. The smaller the fragment, the less immunogenic it may be. This has resulted in Fab'2, Fab fragments, to peptide-like molecules comprising only one of the three CDR's present on each heavy and light chain of  
25 mammalian antibodies, i.e. CDR3 of the heavy chain. In genetically engineering antibodies ways have also been found to avoid having to express at least two different subunits (a heavy and a light chain) for obtaining one antibody. This has been achieved by making a fusion through a  
30 peptide linker between a heavy and a light chain subunit. This results in so-called single chain antibodies, which will be discussed in more detail later on.

Apart from the drawbacks related to immunogenicity of  
35 monoclonal antibodies, there is also a limitation in the specificities obtainable with monoclonal antibodies. Nature

employs a specific process in making CDR's of antibodies through a complex process of hypermutations, rearrangements and insertions in the gene fragment encoding the CDR, followed by a selection of antibody producing cells in primary and secondary lymphoid organs. The result of these processes is that the variety of binding specificities that can be produced in an animal is limited. As one result thereof monoclonal antibodies are not always useful in the application they are intended for, such as for example immunohistochemistry.

Immunohistochemical and immunofluorescent procedures involving tissue sections generally consist of the steps of embedding a tissue in an embedding medium, with or without prior or subsequent fixation or freezing of the tissue, cutting the embedded tissue in thin sections, incubating the tissue sections with monoclonal antibodies, amplification of the signal and visualisation of cells that have bound antibody by an enzymatic reaction and precipitation of substrate or by fluorescent labels.

A drawback of conventional monoclonal antibodies is that they often do not bind to target epitopes in fixed tissue sections, presumably as a result of destruction of the target epitope by the fixation procedure. Fixation and embedding procedures such as paraffin embedding and formalin fixation, that are optimal for preservation of the anatomical structure of the tissue and are routinely used for medical diagnosis, lead to extensive protein denaturation thus limiting the application of monoclonal antibodies that often bind to native epitopes.

Another breakthrough in specific binding technology involves the use of replicable genetic display packages. The term replicable genetic display package or display package describes a biological particle which has genetic information providing the particle with the ability to replicate. The package can display a fusion protein including an antibody,

antibody fragment, protein or peptide. The fusion protein is presented by the display package in a context which permits the antibody to bind a target structure that is contacted with the display package. The display package can be, for example, derived from bacterial cells, bacterial spores or bacterial viruses. In phage display technology a gene fusion is made between a sequence encoding a peptide of interest and a sequence encoding a protein displayed on the outside of a phage (actually the peptide sequence is inserted in the proteins encoding sequence). Libraries of very large numbers of display packages containing all kinds of different peptides within one of their surface proteins can be easily produced and selected and tested based on their binding affinity for other molecules or cells, etc.

Parts of human antibody genes and sequences encoding single chain variable regions have been inserted in such phage genes and thus antibody fragment phage display libraries have been obtained. Display libraries of antibody variable regions may be constructed by a variety of methods. In general, a large collection of antibodies or antibody-fragments is expressed on the surface of a replicable genetic display package. In an exemplary embodiment of the present invention, the display package is a phage particle which comprises an antibody-coat protein fusion; the nucleotide sequences used for construction of the antibodies are of human origin. The library of surface expressed antibody molecules is brought into contact with the target antigen for affinity selection of antibodies that bind to the antigen. This procedure allows the rapid screening of very large collections of displayed antibody molecules, yields human antibodies and does not require immunization of an animal. Novel specificities may be obtained from such libraries.

Phages expressing antibody specificities of interest are selected from libraries in a four step procedure 1) affinity

binding of phages to the target antigen, 2) removal of non-bound phages by washing 3) elution of bound phages and 4) infection and propagation of eluted phages in E Coli bacteria. A variety of targets has been successfully employed to obtain phage antibodies from libraries by affinity selection. Phage antibody affinity selections have been performed on purified antigens coated to a solid phase, on intact eukaryotic cells in suspension or in monolayers and on prokaryotic cells.

The language "affinity binding" and "affinity selection" as well as "differential binding" refer to the separation of members of the peptide or antibody display library based on the differing abilities of these molecules on the surface of each of the display packages of the library to bind to the target structure. Examples of affinity selection include affinity chromatography, immunoprecipitation, fluorescence activated cell sorting, agglutination and plaque lifts.

In all phage antibody affinity selection procedures, it is imperative to obtain a large collection of binding phages in the first round of selection, preferably binding to diverse regions of the target. In subsequent rounds of selection, procedures may be implemented to select from this collection those antibodies that fulfill specific requirements.

The present invention provides further selection methods whereby novel specificities can be easily obtained and/or useful antibody fragments and epitopes (or even any combination of two peptides or proteinaceous substances having binding affinity for each other) can be identified and obtained.

In one embodiment the invention provides a method for identifying a peptide or antibody capable of specific binding to a proteinaceous target, comprising displaying the peptide or antibody on the surface of a replicable display package,

synthesizing sets of oligopeptides derived from the proteinaceous target on a solid phase and contacting the specific binding peptide with the solid-phase-bound oligopeptide to allow for binding. The peptides expressed on the replicable display package may be linear or have another conformation as for example in di-sulfide bridged circular peptides or peptides that contain more than one disulfide bridge and have an unpredictable conformation. In addition, the target oligopeptides may be linear or comprise another configuration.

Thus the present invention provides a method that allows the directed pre-selection of phage antibodies (or other binding peptides) to defined sets of oligopeptides representing (sub)regions of the target molecule. For the purpose of the present invention, the term antibody in its various grammatical forms is art-recognized and includes immunoglobulin molecules and immunologically active portions and/or derivatives thereof, i.e. molecules that contain an antigen binding site. The peptides are synthesized in overlapping or non-overlapping sets of multimers on polystyrene rods or other solid phases such as membranes or beads. In the preferred embodiment, polystyrene rods used for pepscan technology are used. After selection on solid phase-bound rods, collections of phages enriched for anti-oligopeptide phage antibodies can be used in subsequent rounds to select anti-oligopeptide phages that also bind to the target molecule in a another configuration or to select antibodies that discriminate between various forms of the target proteinaceous substance. Many examples of anti-oligopeptide antibodies that bind to the target protein the oligopeptide is derived from have been described in the scientific literature. Herein oligopeptides are defined as comprising 6-30, preferably 8-20 amino acid residues. They will also be simply referred to as peptides.

The term pepscan relates to a procedure for rapid concurrent synthesis on solid supports of hundreds of peptides, essentially as described (Geysen H., et al., Proc. Natl. Acad. Sci. USA 81, 3998 (1984)). In the original procedure, the peptides are synthesized on polyethylene rods as solid support. Other solid supports such as membraneous filters or beads may also be used. Various peptide formats may be synthesized including linear peptides and di-sulfide bridged circular peptides.

Thus in one embodiment the present invention provides a method for generating antibodies specific for sets of peptides synthesized on a solid phase such as the polyethylene rods used in pepscan technology. The subject method generally comprises the steps of synthesizing on a solid phase sets of peptides spanning a target molecule or a subregion thereof; the construction of a display library of antibodies or peptides; the selection of displayed peptides or antibodies that bind to the solid-phase-bound peptides using affinity selection and affinity absorption techniques.

In exemplary embodiments, the display library of peptides or antibodies can be a phage display library or a display library generated on bacterial or yeast cell surface. After one or multiple rounds of selection, the binding displayed antibodies or peptides may be subjected to additional rounds of selection on peptides or on the target protein in its native or denatured configuration. In exemplary embodiments, the subject method can be used to isolate anti-peptide antibodies that also bind to the target protein in its native configuration, for example a marker expressed on normal or malignant eukaryotic cells, or in denatured configuration such as markers expressed on cells in fixed tissue sections. Likewise, the subject method can be used to generate antibodies which can discriminate between a protein and other related forms of the protein, by using a

subtractive approach. The related forms of the protein can differ by one or more amino acids from the target protein or differ as a result of post translational modifications such as glycosylation.

5

In an illustrative embodiment of the subject method, a specific antibody can be generated by affinity selection using a synthetic antibody phage display library. The phage library is incubated with the pepscan block containing rods covered with peptides spanning the extracellular region of a transmembrane protein and phages are allowed to bind to individual rods. After removal of non-bound phages by washing, bound phages are eluted from the rods and propagated in bacterial cells. Multiple subsequent rounds of selection may be performed using the same approach. Alternatively, the binding phages from the first round of selection on solid phase-bound peptides may be selected for binding to the protein in another configuration such as that expressed on a cell or a recombinant protein. In the subtractive approach, phages eluted from the peptides in the first and/or subsequent round(s) of selection may be incubated with rods covered with other peptides, such as peptides differing one or more amino acids from the peptide used for the initial selection, in order to absorb phages that crossreact with related molecules.

25

It is contemplated that the present invention can be applied to obtain bi-specific antibodies that bind to two non-overlapping epitopes on the same molecule of mono meric antigen. Such antibodies have been shown to have enhanced affinity due to the chelate effect (Cheong, H.S., et al. Biochem. Biophys. Res. Comm 173, 795 (1990)). The DNA encoding the pool of phage antibodies affinity selected for binding to a set of overlapping peptides may be recloned into vectors permitting the display of various formats of bi-specific antibodies. The displayed bi-specific antibodies may be affinity selected for binding to the target protein to

30

35

obtain high affinity binding phages. Using the same strategy, bi-specific antibodies may be obtained that bind to two epitopes that are expressed on different molecules.

5 It is also contemplated by the present invention that individual antibodies or peptides, and the genes or oligonucleotides encoding these antibodies or peptides, can be isolated from the phage display library after affinity enrichment. These genes and oligonucleotides can be used for  
10 subcloning into expression vectors for production of antibodies and derivatives of the oligonucleotides can be used for nucleotide sequencing and the information used to produce chemically synthesized peptides or peptides produced by recombinant DNA methods such as production in prokaryotic  
15 or eukaryotic expression systems. In an alternative embodiment the present invention provides a method for identifying a peptide capable of specific binding to a fixated biological target, comprising displaying the peptide on the surface of a replicable display package and  
20 contacting said specific binding peptide with the fixated target to allow for binding.

Thus the present invention provides a method that facilitates the selection of antibodies that are suitable for  
25 immunohistochemical and immunofluorescent procedures with tissue sections that have been embedded and fixed with a variety of procedures. In addition, the procedure facilitates the isolation of antibody specificities that bind to particular anatomical regions of a tissue section, for  
30 example a region infiltrated by tumor cells by using a subtraction procedure and/or micromanipulation of individual cells or small groups of cells. The method employs large libraries of antibodies or peptides expressed on the surface of a replicable display package. The display libraries may be  
35 constructed by a variety of methods. In an exemplary embodiment the display package is a phage particle which



comprises an antibody-coat protein fusion; the nucleotide sequences used for construction of the antibodies are partially synthesized and assembled in vitro. These libraries contain many novel specificities, including antibodies  
5 against self antigens.

In the basic approach, phages expressing antibody specificities of interest are selected from this library in again a four step procedure 1) binding of phages to the  
10 embedded and fixed tissue sections, 2) removal of non-bound phages by washing 3) elution of bound phages and 4) infection and propagation of eluted phages in *E. Coli* bacteria. To obtain specific phage antibodies that bind to a particular tissue or cell type, the method may be extended by absorbing  
15 eluted phage antibodies to remove unwanted specificities. This may be achieved by incubating eluted phage antibodies with sections of other tissues or with sections of the same tissue not containing the target cell of interest and subsequent propagation of unbound phages. In addition, to  
20 obtain phage antibodies specific for a particular cell type, for example defined by anatomical localization, individual or groups of cells or pieces of tissue and attached phages may be scraped from the tissue section by a micromanipulator.

25 The present invention makes available a powerful directed approach for isolating specific antibodies or peptides by antibody or peptide display technologies in combination with solid-phase-dependent peptide synthesis.

30 The present invention further makes available a powerful directed approach for isolating specific antibodies or peptides by antibody or peptide display technologies in combination with selection procedures on tissue sections or whole tissues.

In one exemplary embodiment of the present invention, the display package is a phage particle which comprises an antibody fusion coat protein that includes the amino acid sequence of an antibody variable region. A library of replicable phage vectors, especially phagemids encoding a library of antibody fusion coat proteins is generated and used to infect suitable host cells. The antibody variable genes used to construct the phage library may be entirely derived from the lymphocytes of individuals or (partly) assembled in vitro as in semi-synthetic phage antibody display libraries. An important determinant of the composition and diversity of these in vitro immune systems is the source of antibody genes used as building blocks to construct the library. Libraries may be assembled from the V regions expressed by the B lymphocytes of an individual known to have mounted a particular immune response as a result of immunization or exposure to an infectious agent. The B cell repertoire in the lymphoid organs of these individuals is enriched for antigen-specific B cells and plasma cells that have undergone clonal expansion and antigen affinity maturation, thus increasing the likelihood of selecting high affinity antibody fragments from the eventual "immunized" phage library. Libraries may be constructed from the V regions expressed by the B cells of "non-immunized" individual in an attempt to recruit all the diversity generated by the natural immune system. Importantly, it should be noted that the B lymphocytes of naive, immunized or infected individuals have been through positive and negative selection forces in primary and secondary lymphoid organs. The net effect of these processes is a B cell repertoire that is less diverse than potentially attainable, based on rearrangement of immunoglobulin gene segments, N region insertions and somatic hypermutation. This in vivo antibody repertoire selection is mirrored in the eventual "naive" phage library. An alternative approach to create diverse libraries exploits the use of large collections of cloned V

genes to which randomized CDR3 and JH regions are fused in vitro by PCR. The diversity of these "semi-synthetic" libraries is not constrained by the forces of selection acting in the natural immune system. It may be anticipated that such libraries contain specificities not present in the actual antibody repertoires of humans. In the exemplary embodiment a synthetic library is used because it is expected to contain the novel anti-peptide specificities to be selected.

Phage particles formed from the chimaeric protein can be separated by affinity selection based on the ability of the phage-associated antibody to bind to the target peptide synthesized on rods of a pepsan block. The rods on the pepsan block are decorated with individual peptides, the amino acid sequence of which is determined by the amino acid sequence of the target protein. After one or multiple rounds of selection on peptides synthesized on the pepsan block, the phages are selected for binding to the target protein in its native or denatured configuration. For example, the target protein may be a purified recombinant protein expressed in eukaryotic or prokaryotic cells, or a recombinant protein expressed on the cell surface of a prokaryotic or eukaryotic cell. Phages binding to the target protein are eluted, propagated and eventually analyzed for binding.

After one or multiple rounds of affinity selection on peptides, a subtractive or absorption step may be included to obtain phages that discriminate between two closely related peptides. For example, phages that discriminate between the closely related peptide represented by the amino acid sequences DLVYKDPARPKI and DLVYKDPYRPKI may be obtained by affinity selection on the amino acid sequence DLVYKDPARPKI, followed by absorption of phages binding to the amino acid sequence DLVYKDPYRPKI. In the latter step, non-discriminative

phages recognizing both peptides are removed and not used for propagation. Antibodies produced by this method can for example be used to distinguish various naturally occurring isoforms of a protein or for immunochemical assays for  
5 detecting cell transformations arising due to mutation of an oncogene or an anti-oncogene.

The terms absorption and subtraction refer to the removal of phage antibodies with unwanted specificities from a  
10 collection of phages obtained after or prior to affinity selection. The absorption step itself is an affinity selection step on a target expected not to bear the epitopes of interest. The target used for absorption/selection may be tissue sections, cell lines recombinant proteins, peptides  
15 etc.

### Examples

The invention now being generally described, it will be more  
20 readily understood by reference to the following examples which are included merely for the illustration of certain aspects and embodiments of the present invention and are not intended to limit the invention.

### 25 I. Materials and Methods

Except where indicated otherwise, recombinant DNA methods and microbiological techniques were carried out using standard procedures such as described for example by Sambrook, J. et  
30 al. Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989) and Ausubel, F.M. et al. Current Protocols in Molecular Biology, John Wiley and Sons, Inc. (1995). Specific protocols for propagation and selection of phage antibodies are described  
35 in de Kruif, J., et al. Methods in Molecular Biology, (1996?).

### Construction of a synthetic phage antibody display library

The synthetic antibody library was constructed essentially as described (de Kruif et al., J. Mol. Biol. 248, 97 (1995)).  
5 Briefly, degenerate oligonucleotides were used to add synthetic CDR3 regions to a collection of 49 previously cloned germline VH genes. Subsequently, these in vitro 'rearranged' VH genes were cloned into a collection of pHEN1  
10 phagemid-derived vectors containing 7 different light chain V regions, fused in frame to the gene encoding the phage minor capsid protein geneIII. Introduction of these constructs into bacteria results, in the presence of helper phage, in the expression of scFv antibody fragments as geneIII fusion  
15 proteins on the surface of bacteriophage.

### Preparation of tissue sections

Fixation, freezing, embedding and sectioning of tissues were  
20 carried out using standard procedures such as for example described in Zeller, R in Current Protocols in Molecular Biology, pp 14.1.1-14.1.8, Ausubel, F.M. et al. Eds, Green Publishing and Wiley Interscience, New York.  
Immunohistochemical and immunofluorescent procedures were  
25 carried out using standard protocols such as for example described in Current Protocols in Immunology, pp 5.8.1-5.8.8, Coligan, J.E., et al. Eds. Wiley Interscience, New York.

### Example I: selection using pepscan

30

#### Pepscan synthesis of CD64 peptides

Pepscan synthesis was carried out essentially as described (Geysen, H.M., et al, Proc. Natl. Acad. Sci. USA, 81, 3998  
35 (1984)). In the exemplary embodiment, the extracellular domain of the CD64 molecule was synthesized on 25

polyethylene rods as adjacent, non-overlapping sets of 12-mer peptides, according to the scheme in Table I.

Peptide-specific phages were isolated by affinity binding to the CD64 peptide-covered rods. The rods were blocked by

- 5 incubation for two hours at room temperature by placing the pepsan block for two hours at room temperature in wells of a 96-well microtiter plate containing 300 microliter 1x blocking buffer per well (2x blocking buffer: 10% ovalbumin/10% horse serum/1% Tween 80 in PBS). The phages
- 10 were were mixed 1:1 (v/v) with 2x blocking buffer and incubated for 30 minutes at room temperature. 250 microliter of the blocked phage solution containing approximately  $10^{12}$  phages were added per well of a 96 well plate and the 25 rods of the pepsan block were submerged in individual wells and
- 15 incubated for 3 hours at room temperature. Non-specific phage antibodies were then removed by washing the rods 10 times in PBS/0.5% Tween 80 and one final time in PBS. Phage antibodies bound to the rods were then eluted by adding 300 microliter of 0.1M diethylamine buffer to each well of a microtiter
- 20 plate and incubation for 5 minutes at room temperature. After collection of the 300 microliter, the elution step was repeated. To 600 microliter of eluted phages, 300 microliter of 1M Tris buffer pH 7.4 was added for neutralization. The 900 microliter eluted phage suspension was added to 4 ml 2TY
- 25 containing 5% glucose and 10 microgram/ml tetracycline. Finally, 5 ml XL-1 blue cells (OD600 of 0.5) were added and the mixture incubated for 30 minutes at 37°C. The cells were centrifuged for 30 minutes at 3000 rpm, the supernatant removed by aspiration and the cells resuspended in 500
- 30 microliter of the remaining medium on two TYE plates containing 5% glucose, 100 microgram/ml ampicillin, 10 microgram/ml tetracycline and 5% glucose and the bacteria grown overnight at 37° C. The colonies were scraped from the bacterial plates and used to rescue phage antibody particles as described (de
- 35 Kruif, J., et al., J. Mol. Biol. 248, 97 (1995)). From the resulting library enriched for peptide-binding phage

antibodies,  $10^{12}$  phage antibodies in blocking buffer were incubated with the same peptide-covered rods and subjected to subsequent rounds of selection using the same procedure.

5 Cleaning of pepscan blocks

Blocks used for ELISA or phage selections were cleaned by submerging the pep scan blocks in disruptbuffer (phosphatebuffer/1%SDS/0.1%beta-mercapto-ethanol, pH 7.2) and  
10 placing the block in a ultrasonic waterbath at 70°C for 1 hour.

Analysis of selected phages in ELISA

15 After selection, preparations of monoclonal or polyclonal scFv antibodies were analyzed for the presence of peptide binding in ELISA using the same pepscan blocks used for selections. ScFv preparations instead of phage preparations  
20 were used because the latter generated a high background in ELISA. All incubations are carried out in 96-well microtiter plates. The cleaned rods were soaked in 300 microliter PBS for 30 minutes at roomtemperature and pre-coated by  
25 incubating each pin for 3 hours at roomtemperature with 150 microliter of blocking buffer (5% ovalbumin/5% horse serum/1% Tween 80, filtered over Whatman 41 paper). The rods were  
30 rinsed twice with PBS/0.05% Tween 80 and incubated for 1 hour at roomtemperature with 150 microliter scFv-containing periplasmic preparations prepared as described in de Kruif et al, J. Biol. Chem. 271, 7630 (1996)) mixed with 150  
35 microliter blocking buffer. The rods were washed twice with PBS/0.05% Tween 80 and incubated for 1 hour with 300 microliter culture supernatant of a hybridoma secreting the anti-Myc tag antibody 9E10 (de Kruif, J., et al, J. Biol. Chem. 271, 7630 (1996)). The rods were washed twice with  
40 PBS/0.05% Tween 80 and incubated for 1 hour with 300 microliter of rabbit anti-mouse immunoglobulins conjugated to

peroxidase, diluted 1:500 in PBS/0.05% Tween 80/1% BSA. Finally, the rods were washed thrice with PBS/0.05% Tween 80 and incubated with ABTS substrate. A representative result of an anti-peptide scFv antibody, affinity-selected after three rounds for binding to peptide # 21 on the CD64 block is shown in figure 2. This scFv antibody specifically bound to peptide #21 used as target for selection but not to other peptide-covered rods in the CD64 pepscan block (figure 2).

10 Selection and analysis of pepscan pre-selected phage antibodies by cell sorting.

Phage populations enriched for binding to peptides on the pepscan block may in the second or subsequent rounds be selected for binding to prokaryotic or eukaryotic cells expressing the target protein employing flow cytometry and cell sorting. The procedure for phage selection on intact cells using this procedure has been described in detail (de Kruif, J., et al. Proc. Natl. Acad. Sci. USA, 92,3938).

20 Approximately  $10^{13}$  phage antibody particles are blocked for 15 minutes in 4 ml 4% milkpowder in PBS (MPBS).  $5 \times 10^6$  target cells were added to the blocked phages and the mixture was slowly rotated overnight at 4°C. The following day, cells were washed twice in 50 ml ice-cold PBS/1% BSA. The pelleted cells were resuspended in 2 ml of ice-cold PBS/1% BSA for cell sorting or stained with fluorochrome-labeled monoclonal antibodies for sorting of a subpopulation of cells. To that end, 25 microliter of fluorochrome-labeled antibodies is added to the pelleted cells and after a 20 minute incubation on ice, cells were washed once with 1% BSA/PBS and resuspended in 2 ml ice-cold PBS/1% BSA. Cell sorting was performed on a FACSVantage.

The procedure for flow cytometric analysis of selected phage antibodies has been described in detail elsewhere (de Kruif, J., et al. Proc. Natl. Acad. Sci. USA, 92,3938). For staining



of cells, 100  $\mu$ l monoclonal phage antibody containing approximately  $10^{11}$  particles was blocked by adding 50  $\mu$ l 4% MPBS for 15 minutes at roomtemperature.  $5 \times 10^5$  leucocytes in 50  $\mu$ l PBS/1% BSA were added and incubated on ice for 1 hour.

- 5 The cells were washed twice in ice-cold PBS/1% BSA. To detect cell-bound phages, the cells were incubated in 10  $\mu$ l of 1/200 diluted sheep anti-M13 polyclonal antibody (Pharmacia, Uppsala, Sweden), washed twice and incubated in 10  $\mu$ l of 20  $\mu$ g/ml PE-labeled donkey anti-sheep polyclonal antibody
- 10 (Jackson ImmunoResearch, West Grove, PA), each for 20 minutes on ice.

Selection of pepscan pre-selected phage antibodies on whole tissues or tissue sections.

15

- Tissue sections from target organs are prepared and fixed by standard methods. For example, tissues may be submerged in OCT embedding medium and gently frozen by dipping in liquid nitrogen for 1 minute. Using a cryostat, 10 micrometer frozen
- 20 sections are prepared and collected on microscope slides. The sections are dried for 10 minutes at roomtemperature and fixed in 100% acetone for 5 seconds, followed by drying at roomtemperature. Before incubation with the phage library, the tissue sections are sealed with wax to prevent spreading
- 25 of fluid and blocked by incubation with MPBS for 30 minutes at roomtemperature. After the blocking step, a total of  $10^{12}$  phages from a phage antibody library pre-selected for binding to peptides is blocked with MPBS in a volume of 100 microliter and added to the droplet of MPBS on the slide. The
- 30 tissue sections are incubated overnight in a moist chamber at 40°C to facilitate binding of phage antibodies.

- The sections are rinsed with a total of 2 liter PBS/0.5% Tween 20 for 2 hours at roomtemperature with ten 200 ml
- 35 changes of PBS/0.5% Tween 20. After rinsing, the phage antibodies are eluted by placing 300 microliter 76 mM citric

acid buffer (pH 2.5) on the slide and incubation for 5 minutes at roomtemperature. Add 150 microliter of 1 M Tris-HCL buffer are added to the slide for neutralization and the suspensi on is brought in 3 ml 2 TY medium in a 15 ml tube. 3 ml of XL-1 blue bacteria is added to the tube and infection and production of phages are allowed to proceed as described in the standard protocol. Afer 2-3 rounds of selection, monoclonal phage antibodies are prepared for analysis.

#### Analysis of selected phage antibodies by immunohistochemistry

Binding of monoclonal phage antibodies to tissues was determined by immunohistological screening. To this end, 5  $\mu$ m frozen tissue sections were collected on gelatin coated microscope slides. Sections were then incubated with 50 microliter phage solutions containing  $10^{10}$  phage particles in a moist chamber for 30 min. Non-adherent phages were removed by vigorously rinsing the slides in M-PBS-Tw. Bound MoPhabs were visualized by incubation of the sections with a 1:50 diluted sheep anti-M13 peroxidase conjugate, followed by incubation with the substrate diamino benzidine.

#### Construction and selection of bi-specific antibodies from collections of phages antibodies affinity selected on pepsan blocks.

Monoclonal phage antibodies selected for binding to the CD64 peptides may be pooled and recloned in display vectors permitting the display of bi-specific antibodies such as for example vectors for diabodies (Holliger, P. et al., Proc. Nat. Acad. Sci. USA, 90, 6444 (1993) or leucine zippered bi-specific antibodies (de Kruif, J., et al. J. Biol. Chem. 271, 7630 (1996).

In the exemplary embodiment, a phagemid vector was constructed that facilitates the in vivo bacterial expression and assembly of bispecific (scFv)<sub>2</sub> fragments as secreted molecules or g3p fusion proteins on the surface of bacteriophages. The Fos and Jun scFv leucine zipper constructs we described (de Kruif, J., et al. J. Biol. Chem. 271, 7630 (1996); figure 3, 1 and 2) are PCR amplified using primers M13R and PDIM3 (5'-TTT GCA TTC AAG CTT TTA TTA GCC CGC ATA GTC AGG AAC ATC GTA TGG GTA TGC GGC AGC GCA ACC ACC). Primer PDIM3 replaces the Myc tag with a haemagglutinin (HA) tag and adds a stopcodon and a HindIII site to these fragments. PCR products are digested with HindIII and cloned into vector pHEN1 (Hoogenboom, H.R., et al., Nucleic Acid Res. 19, 4133 (1991)) containing a scFv gene fused to the complementary leucine zipper and a Myc tag. In the resulting constructs both scFv-zipper fragments are encoded by a single transcript. Two ribosome binding sites allow the proteins to be translated individually, the first scFv fragment fused to a Fos zipper domain and an HA tag, the second scFv to a Jun zipper and a Myc tag (Figure 3, 3). To determine the effect of the position of the scFv-zippers relative to the g3p protein, an additional construct was made in which the scFv-zipper fragments were reversed (Figure 3, 4). In suppressor E. coli strains, an amber codon inserted between the Myc tag and gene 3 permits the production of phage particles displaying a scFv fragment fused via a zipper region to gene 3. The second scFv will associate with the phage in the periplasmic space by heterodimerization of the Fos and Jun leucine zippers, thus creating a phage expressing a bispecific antibody on its surface. In non-suppressor strains, bispecific (scFv)<sub>2</sub> fragments linked by a Fos Jun leucine zipper will be assembled in the periplasmic space.

Expression of (scFv)<sub>2</sub> fragments in non-suppressor bacterial strain SF110-F<sup>+</sup> was monitored by SDS-PAGE followed by

blotting to nitrocellulose and detection with Myc tag- and HA tag-specific monoclonal antibodies (MoAbs) (Figure 4, lower panel). In reducing SDS-PAGE, constructs 3 and 4 show expression of HA and Myc tagged proteins of the predicted size. The expression level of Myc tagged scFv fragments obtained with constructs 3 and 4 is lower when compared to that obtained with constructs 1 and 2, perhaps as a result of the simultaneous expression of a second scFv in bacteria harboring the former constructs. In non-reducing SDS-PAGE, only proteins approximately twice the size of a scFv-zipper are visible in all lanes (Figure 4, upper panel). The anti HA-MoAb detects multiple bands, whereas the anti Myc-MoAb detects only one. One of the fragments visualized with the anti-HA MoAb migrates at the same position as the fragments visualized with the anti-Myc MoAb. We hypothesize that this band represents the bispecific antibody and that the other HA-tagged proteins are homodimers resulting from a higher expression level of the first scFv-zipper in these constructs.

Functionality of the (scFv)<sub>2</sub> fragments was analysed in an ELISA (Figure 5). Using both the Myc tag- and HA tag-specific Moabs for detection, specific binding of the (scFv)<sub>2</sub> fragments to both target antigens could be demonstrated. In a sandwich ELISA we show the (scFv)<sub>2</sub> fragments are truly bispecific in linking two different antigens. Recombinant phage antibody particles were produced using constructs 1-4 and analysed in an ELISA (Figure 5). Phages from constructs 2 and 3, containing the Jun-zipper fused directly to gene 3 show some non-specific binding to irrelevant antigens. Phages produced using constructs 1 and 4, containing the Fos-zipper fused to gene 3 recognize their target antigens only.

To test the feasibility of selecting bispecific phage antibodies from a phage repertoire, bispecific phage

particles from construct 4 were spiked in a 1:1000 ratio in an antibody phage display library and subjected to one round of selection in immunotubes coated with either IgG or DNP. The frequency of bispecific phages after selection was  
5 estimated by PCR on individual colonies using primers M13R and FOSCON (5'-CGC CAG GAT GAA CTC C, situated in the Fos zipper). Frequencies of bispecific phages after selection were 0/32 (spiked library), 9/32 (selected on IgG) and 6/32 (selected on DNP). Calculated enrichment factors ( $\pm 250x$ ) are  
10 comparable to those obtained using monospecific phage antibodies.

This vector permits the expression of bispecific antibodies that can be functionally expressed on phages using Fos and  
15 Jun leucine zippers. It is therefore possible to construct libraries of phages expressing two different scFv fragments and select the desired combinations from these. Furthermore, selected bispecific (scFv)<sub>2</sub> fragments can be produced in non-suppressor strains and purified using anti-Myc followed  
20 by anti-HA affinity columns.

For the selection of bi-specific antibodies from phage antibody collections pre-selected for binding to peptides on  
25 pepscan blocks, care is taken to restrict the number of selection rounds (usually 2-3) on the rod-bound peptides to ensure enrichment for binders but to prevent loss of diversity by overselection. The V gene pools encoding the phage antibodies enriched for binding to the peptides are recloned into the leucine-zipper constructs and bi-specific  
30 phage antibodies binding to the target protein are selected by any of the afore-described methods. In addition, it may be envisaged that higher affinity bi-specific antibodies are directly isolated by performing phage selection with the aid of surface plasmon resonance (Jonsson, U., et al.  
35 Biotechniques, 11: 620 (1991)).

## Example 2: selection on sections

## Preparation of tissue sections

Tissue sections from target organs are prepared and fixed by standard methods. For example, tissues may be embedded in an embedding compound such as OCT and gently frozen by dipping in liquid nitrogen for 1 minute. Using a cryostat, 10 micrometer frozen sections are prepared and collected on microscope slides. The sections are dried for 10 minutes at roomtemperature and fixed in 100% acetone for 5 seconds, followed by drying at roomtemperature.

Alternatively, tissues may be fixed by incubation for several hours in 4% freshly prepared paraformaldehyde solution, dehydrated by serial passage through increasing concentrations of ethanol and embedding in paraffin wax. Paraffin blocks are cut into 6-8 micrometer sections. Instead of tissue sections, freshly isolated or cultured cells may be used and subjected to the same procedures.

Affinity selection of phages on tissue sections

The frozen tissue sections are briefly dried at room temperature. The paraffin-embedded tissue sections are placed at 60°C for 1 hour and deparaffinized by immersing in three changes of fresh xylene, 4 minutes each, followed by two changes of 100% ethanol, 2 minutes each. Tissue sections are rehydrated in solutions of decreasing concentrations of ethanol and finally water, and washed with PBS.

Before incubation with the phage library, the tissue sections are sealed with wax to prevent spreading of fluid over the object glass, and blocked by incubation with 1% non-fat milkpowder in phosphate buffered saline (MPBS) for 30 minutes at roomtemperature. After the blocking step, a total of  $10^{13}$  phages from the synthetic phage antibody display library is

blocked with MPBS in a volume of 300 microliter and added to the droplet of MPBS on the slide. The tissue sections are incubated over night in a moist chamber at 40°C to facilitate binding of phage antibodies.

5

The sections are rinsed with a total of 2 liter PBS/0.5% Tween 20 for 2 hours at roomtemperature with 10 200 ml changes of PBS/0.5% Tween 20. After rinsing, the phage antibodies are eluted by placing 300 microliter 76 mM citric acid buffer (pH 2.5) on the slide and incubation for 5 10 minutes at roomtemperature. 150 microliter of 1 M Tris-HCL buffer is added to the slide for neutrlaization and the suspension is brought in 3 ml 2 TY medium in a 15 ml tube. 3 ml of XL-1 blue bacteria are added to the tube and infection 15 and production of phages are allowed to proceed as described in the standard protocol. Afer 2-3 rounds of selection, monoclonal phage antibodies are prepared for analysis by immunohistochemistry.

20 Analysis of selected phage antibodies by immunohistochemistry

Binding of monoclonal phage antibodies to tissues was determined by immunohistological screening. To this end, 5 µm frozen tissue sections were collected on gelatin coated 25 microscope slides. Sections were then incubated with 50 microliter phage solutions containing approximately 10<sup>10</sup> phage particles in a moist chamber for 30 min. Non-adherent phages were removed by vigorously rinsing the slides in MPBS-Tw. Bound phage antibodies were visualized by incubation 30 of the sections with a 1:50 diluted sheep anti-M13 peroxidase conjugate, followed by incubation with diaminobenzidine substrate.

Absorption of selected phages antibodies for unwanted specificities

A collection of phage antibodies selected for binding to a tissue section may be absorbed for unwanted phage antibody specificities. After elution and neutralization of bound phage antibodies, a 4% (w/v) MPBS solution is added to the eluate to reach a 1% concentration of non-fat milk. The phage solution is subsequently incubated with a tissue sections of for example another tissue or a different anatomical region of the same tissue. Non-binding phages are removed and propagated directly or incubated with another tissue section for another round of absorption. This absorption procedure may be repeated multiple times before the eventual non-binding phages are propagated in bacteria.

Isolation of specific phage antibodies from tissue sections using a micromanipulator.

Individual cells or small regions of tissue and binding phages may be scraped from tissue sections using a micromanipulator or similar device, essentially as described (Kupper, R., et al. EMBO-J, 12 4955 (1993)). This procedure allows the isolation of phages that bind to particular anatomical structure or individual cells in a tissue section. For example, this procedure allows the isolation of endothelial of bloodvessels penetrating a tumor mass or allow the isolation of individual tumor cells. Once scraped from the tissue section, the bound phages are eluted and propagated as described in the basic protocol.



Legends to the figures

## Figure 1.

General principle of affinity selection of genetic display packages by binding to peptides synthesized on the rods of a pepsan block.

## Figure 2.

ELISA results of a phage antibody affinity selected for binding to peptide #21 on the CD64 pepsan block. No binding above background to other peptides on the block was observed.

## Figure 3.

Starting material (1 and 2) and bispecific constructs (3 and 4). LacZ promoters (triangles), ribosome binding sites (circles) pelB leaders (P), Fos-zippers (F), Jun-zippers (J), Myc tag (M), HA tag (H) and stopcodons (arrows) are indicated. Closed boxes and hatched boxes represent different scFv specificities.

## Figure 4.

SDS-PAGE / Western blotting analysis. Periplasmic preparations from expressed constructs 1-4 were run under non-reducing (top panel) and reducing (bottom panel) conditions. Proteins were visualized using an HA-tag specific MoAb (left lanes) and a Myc-tag specific MoAb (right lanes) followed by a peroxidase- conjugated antibody.

## Figure 5.

ELISA analysis using construct 1-4 derived antibody fragments. In standard ELISA's, (scFv)<sub>2</sub> fragments were allowed to bind to antigen coated wells and detected using anti-Myc and anti-HA MoAbs followed by a peroxidase-conjugated antibody. Sandwich ELISA's using bispecific (scFv)<sub>2</sub> fragments were performed as described (de Kruif, J et al. J. Biol. Chem. 271, 7630). In phage ELISA,

phage antibodies were allowed to bind to antigen coated wells and developed using a peroxidase conjugated anti-M13 antibody. Numbers represent absorbance values. Values three times higher than the background binding to milk-blocked  
5 wells are considered to be positive (gray boxes).

	pin #	amino acid sequence	pin #	amino acid sequence
	1	MWFLTLLLLWVP	14	KTNISHNGTYHC
	2	VDGQVDTTKAVI	15	SGMGKHRYTSAG
5	3	SLQPPWVSVFQE	16	ISVTVKELFPAP
	4	ETVTLHCEVLHL	17	VLNASVTSPLE
	5	PGSSSTQWFLNG	18	GNLVTLSCETKL
	6	TATQTSTPSYRI	19	LLQRPGLQLYFS
	7	TSASVNDSGEYR	20	FYMGSKTLRGRN
10	8	CQRLSGRSDPI	21	TSSEYQILTARR
	9	QLEIHRGWLLQ	22	EDSGLYQCEAAT
	10	VSSRVFTEGEPL	23	EDGNVLKRSPEL
	11	ALRCHAWKDKLV	24	ELQVLGLQLPTP
	12	YNVLYYRNGKAF	25	VWFHVLFLAVG
15	13	KFFHWNSNLTL		

Table 1. Sets of oligopeptides representing the extracellular domain of CD64.

CLAIMS

1. A method for identifying a peptide capable of specific binding to a proteinaceous target, comprising displaying the peptide on the surface of a replicable display package, synthesizing oligopeptides derived from the proteinaceous  
5 target on a solid phase and contacting the specific binding peptide with the oligopeptide to allow for binding.
2. A method for identifying a peptide capable of specific binding to a fixated biological target, comprising displaying the peptide on the surface of a replicable display package  
10 and contacting said specific binding peptide with the fixated target to allow for binding.
3. A method for distinguishing between peptides capable of specific binding to a proteinaceous antigen and peptides not having that capability comprising displaying candidate  
15 peptides on the surface of a replicable display package, synthesizing oligopeptides derived from the proteinaceous antigen on a solid phase and contacting the candidate peptides with the oligopeptides to allow for binding and washing the solid phase to remove the display packages not  
20 specifically bound.
4. A method for distinguishing between peptides capable of specific binding to a fixated biological target and peptides not having that capability comprising displaying candidate peptides on the surface of a replicable display package,  
25 contacting said specific binding peptide with the fixated target to allow for binding and washing the fixated biological target to remove the display packages not specifically bound.
5. A method according to anyone of the afore going claims,  
30 whereby the replicable display package is a phage particle.
6. A method according to anyone of the afore going claims, whereby the replicable display package is a bacterium, a yeast or a spore of a microorganism.

7. A method according to claim 5, whereby the specific binding peptide is displayed on the surface of the phage by insertion of its encoding sequence in a gene encoding a surface protein of said phage.

5 8. A method according to anyone of the foregoing claims, whereby the displayed peptide is an immunoglobulin heavy chain, an immunoglobulin light chain, a heavy-light chain pair, a VH, a VL, a Fab, a Fv, an scFv or a di-sulfide-bridged Fv.

10 9. A method according to anyone of the foregoing claims whereby the specific binding peptide is a single chain antibody fragment, preferably an ScFv.

10. A method according to anyone of the foregoing claims, further comprising a step whereby the displayed peptides are  
15 contacted with a sample not containing the target of interest.

11. A method for screening a library of replicable display packages for peptides capable of specific binding to a proteinaceous target or a fixated biological target comprising  
20 subjecting the peptides in the library to a method according to any one of the foregoing claims.

12. A method according to claim 11 wherein said library is a phage display library.






Figure 1





# CD64 pepscan using scFv clone F2




Figure 2






Figure 3







Figure 4



	1	2	3	4	
<i>IgG</i>	2.11	0.09	0.61	0.26	<i>(scFv)<sub>2i</sub>; α-Myc</i>
<i>DNP</i>	0.07	2.63	1.74	1.22	
<i>Milk</i>	0.06	0.08	0.06	0.06	
<i>IgG</i>	0.07	0.07	1.62	0.56	<i>(scFv)<sub>2i</sub>; α-HA</i>
<i>DNP</i>	0.07	0.06	1.95	2.26	
<i>Milk</i>	0.06	0.06	0.06	0.06	
<i>IgG + DNP</i>	0.06	0.08	0.79	0.51	<i>(scFv)<sub>2i</sub>; sandwich</i>
<i>IgG</i>	1.56	0.55	0.96	1.05	<i>biphab; α-M13</i>
<i>DNP</i>	0.09	1.81	1.62	1.19	
<i>Milk</i>	0.06	0.62	0.35	0.09	

Figure 5





A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 G01N33/554 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**BEST AVAILABLE COPY**

C. DOCUMENTS CONSIDERED TO BE RÉLEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 15677 A (PROTEIN ENGINEERING CORP) 17 September 1992. see the whole document	1,3,5-7, 11,12
X	DE KRUIF ET AL.: "Selection and application of human single chain Fv antibody ...." JOURNAL OF MOLECULAR BIOLOGY, vol. 248, no. 1, 21 April 1995, LONDON, pages 97-105, XP000196590 cited in the application see the whole document	1,3,5, 8-12
Y		2,4,6,7
Y	WO 92 08738 A (ABBOTT LAB) 29 May 1992 see page 8, line 21 - line 23	2,4
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

2 December 1997

Date of mailing of the international search report

19.12.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Ceder, 0

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 15982 A (GENZYME CORP) 15 June 1995 see claims 1-5,65 ---	6
Y	WO 95 31723 A (FRYKBERG LARS ; JACOBSSON KARIN (SE)) 23 November 1995 see abstract ---	7
A	GEYSEN ET AL: "USE OF PEPTIDE SYNTHESIS TO PROBE VIRAL ANTIGENS FOR EPITOPES TO A RESOLUTION OF A SINGLE AMINO ACID" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 81, July 1984, WASHINGTON DC, pages 3998-4002, XP000609021 cited in the application see abstract -----	1,3

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Patent Application No

PCT/NL 97/00557

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9215677 A	17-09-92	US 5223409 A	29-06-93
		JP 7501923 T	02-03-95
		AU 1545692 A	06-10-92
		AU 1578792 A	06-10-92
		AU 1581692 A	06-10-92
		AU 8740491 A	28-04-92
		CA 2105300 A	02-09-92
		CA 2105303 A	02-09-92
		CA 2105304 A	02-09-92
		EP 0575485 A	29-12-93
		EP 0573603 A	15-12-93
		EP 0573611 A	15-12-93
		JP 6510522 T	24-11-94
		JP 7501203 T	09-02-95
		US 5403484 A	04-04-95
		US 5571698 A	05-11-96
		WO 9206191 A	16-04-92
		WO 9215605 A	17-09-92
		WO 9215679 A	17-09-92
		US 5663143 A	02-09-97
WO 9208738 A	29-05-92	AU 662503 B	07-09-95
		AU 9077791 A	11-06-92
		CA 2095735 A	08-05-92
		EP 0556320 A	25-08-93
		JP 6502076 T	10-03-94
WO 9515982 A	15-06-95	AU 1432195 A	27-06-95
		EP 0733070 A	25-09-96
		JP 9506262 T	24-06-97
		CA 2175482 A	15-06-95
WO 9531723 A	23-11-95	AU 2541595 A	05-12-95
		EP 0760100 A	05-03-97

